

STUDY OF VENOUS TONUS IN THE FOREARM DURING
SIMULATED WEIGHTLESSNESS

D. Kaiser and O. H. Gauer


(NASA-TT-F-15228) STUDY OF VENOUS TONUS
IN THE FOREARM DURING SIMULATED
WEIGHTLESSNESS (Kanner (Leo) Associates)
3 p HC \$3.00 CSCL 06P

N74-14828

Unclass
27353

G3/04

Translation of "Untersuchungen über den Venentonus am Unterarm
bei simulierter Schwerelosigkeit," Pflugers Archiv (European
Journal of Physiology), Vol. 289, 1966, pp. R76-77



STANDARD TITLE PAGE

1. Report No. NASA TT F-15,228		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle STUDY OF VENOUS TONUS IN THE FOREARM DURING SIMULATED WEIGHTLESSNESS				5. Report Date January 1974	
				6. Performing Organization Code	
7. Author(s) D. Kaiser and O. H. Gauer, Physiologisches Institut der Freien Universität, Berlin				8. Performing Organization Report No.	
				10. Work Unit No.	
9. Performing Organization Name and Address Leo Kanner Associates Redwood City, California 94063				11. Contract or Grant No. NASW-2481	
				13. Type of Report and Period Covered Translation	
12. Sponsoring Agency Name and Address National Aeronautics and Space Adminis- tration, Washington, D.C. 20546				14. Sponsoring Agency Code	
15. Supplementary Notes Translation of "Untersuchungen über den Venentonus am Unterarm bei simulierter Schwerelosigkeit," Pflugers Archiv (European Journal of Physiology), Vol. 289, 1966, pp. R76-77					
16. Abstract A reduction in venous tonus, assumed to be partly a function of temperature, is observed directly after immersion in a thermally neutral bath. Several hours of immersion causes further relaxation of veins.					
17. Key Words (Selected by Author(s))			18. Distribution Statement Unclassified-Unlimited		
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 4	22. Price		

STUDY OF VENOUS TONUS IN THE FOREARM DURING
SIMULATED WEIGHTLESSNESS

D. Kaiser and O. H. Gauer,
Physiologisches Institut der Freien Universität, Berlin

After several hours of simulated weightlessness (immersion /R76*
in a thermally neutral bath), a tendency toward collapse occurs
under orthostatic loads (tilting table). In earlier experiments,
it had been possible to show that under these conditions -- after
8 hours of immersion -- plasma volume is reduced by 14% [1]. A
study was now made as to whether a reflex slackening of the veins
also favors the tendency toward collapse. Prior to immersion --
and at regular intervals after immersion -- circumferential
pressure diagrams were taken with the aid of a venous inflatable
cuff, a Whitney gauge [2] and Statham strain gauge manometer about
the upper arm. A reduction in venous tonus, probably a function,
in part, of temperature, occurred immediately after immersion.
Several hours of immersion also usually causes further relaxation
of the veins.

* Numbers in the margin indicate pagination in the foreign text.

REFERENCES

/R77

1. Gauer, O. H., Eckert, P., Kaiser, D., and Linkenbach, H.-J.,
2nd Internat. Symp. on Basic Environmental Problems of Man
in Space, Paris 1965, Vienna, Spring (in press).
2. Whitney, R. J., J. Physiol. (Lond.) 121, 1 (1953).